

Suppression of Entropy Perturbations in Multi-Field Inflation on the Brane

P.R. Ashcroft, C. van de Bruck and A.-C. Davis

*Department of Applied Mathematics and Theoretical Physics, Center for Mathematical Sciences,
University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, U.K.*

(August 2002)

At energies higher than the brane tension, the dynamics of a scalar field rolling down a potential are modified relative to the predictions of General Relativity. The modifications imply, among other things, that steeper potentials can be used to drive an epoch of slow-roll inflation. We investigate the evolution of entropy and adiabatic modes during inflation driven by two scalar fields confined on the brane. We show that the amount of entropy perturbations produced during inflation is suppressed compared to the predictions made by General Relativity. As a consequence, the initial conditions do not matter in multiple field inflation in brane worlds if inflation is driven at energies much higher than the brane tension.

DAMTP-2002-107

Inflation provides an explanation for the origin of structures in the universe. In its simplest form, inflation is driven by a single scalar field, called the inflaton, which slowly rolls down a potential. Perturbations in the inflaton field are adiabatic and obey Gaussian statistics [1]. However, this picture might be over simplistic as there is no reason to believe that inflation is driven solely by a single scalar field. Instead, the possibility that inflation is driven by a system of several scalar fields must be considered. In particular, perturbations are, in general, no longer purely adiabatic. Entropy (or isocurvature) perturbations can be generated. In addition, non-Gaussian fluctuations are now possible. In the case of multiple field inflation, there are no unique background trajectories for the fields and therefore predictions of density perturbations could depend on initial conditions [2].

Current data do not exclude an additional entropy mode [3,4]. Future experiments such as the Microwave Anisotropy Probe (MAP), the Planck Surveyor, the 2 Degree Field Galaxy Survey (2dF) and Sloan Digital Sky Survey (SLOAN) will constrain deviations from the standard predictions of single scalar field inflation. These prospects motivated recent work to consider multi-field inflation and the consequences for density perturbations [5–7].

In the case of brane worlds based on the model of Randall and Sundrum [8], inflation driven by a scalar field confined on the brane has been considered (see e.g. [9–12]). It was shown that the correction of Einstein's equation at high energies lead to a suppression of the usual slow roll parameters due to the enhanced damping of the scalar field. This in turn implies that the spectral index of the spectrum of perturbations is nearer one for a given potential compared to the predictions of General

Relativity. In this letter we point out that the enhanced damping implies that the generation of entropy perturbations is suppressed in the case of multiple-field inflation on the brane for a given potential. We also deduce that steeper potentials are needed to create the same amount of entropy modes as in usual four-dimensional gravity. Furthermore, any transfer of non-Gaussianity from the entropy modes to the adiabatic modes is suppressed. If inflation is driven at energies much higher than the brane tension, the results of density perturbations in multiple field inflation become independent of the initial conditions.

General Relativity: Before we consider inflation on a brane, we review briefly the results obtained in General Relativity [13]. It turns out the form of the equations are not modified in the case of the brane world model we consider.

To be precise, we consider the case of two scalar fields, ϕ and χ , which are slowly rolling down some potential $V(\phi, \chi)$. The equations of motion are governed by the Friedmann equation and Klein-Gordon equations:

$$H^2 = \frac{8\pi G}{3}\rho, \quad (1)$$

$$\ddot{\phi} + 3H\dot{\phi} + V_\phi = 0, \quad (2)$$

$$\ddot{\chi} + 3H\dot{\chi} + V_\chi = 0, \quad (3)$$

where ρ is the energy density of *both* fields. We assume that the fields are slow-rolling, implying that the parameters

$$\begin{aligned} \epsilon_\phi &\equiv \frac{1}{16\pi G} \left(\frac{V_\phi}{V} \right)^2, \quad \epsilon_\chi \equiv \frac{1}{16\pi G} \left(\frac{V_\chi}{V} \right)^2 \\ \eta_{\phi\phi} &\equiv \frac{1}{8\pi G} \left(\frac{V_{\phi\phi}}{V} \right), \quad \eta_{\phi\chi} \equiv \frac{1}{8\pi G} \left(\frac{V_{\phi\chi}}{V} \right) \\ \eta_{\chi\chi} &\equiv \frac{1}{8\pi G} \left(\frac{V_{\chi\chi}}{V} \right) \end{aligned} \quad (4)$$

are small. A convenient formalism to study entropy perturbations was presented in [5] which we will use here. Instead of working with the fields ϕ and χ it is useful to perform a field rotation as follows:

$$\delta\sigma = (\cos\theta)\delta\phi + (\sin\theta)\delta\chi \quad (5)$$

$$\delta s = -(\sin\theta)\delta\phi + (\cos\theta)\delta\chi, \quad (6)$$

with

$$\cos\theta = \frac{\dot{\phi}}{\sqrt{\dot{\phi}^2 + \dot{\chi}^2}}, \quad \sin\theta = \frac{\dot{\chi}}{\sqrt{\dot{\phi}^2 + \dot{\chi}^2}}. \quad (7)$$

σ is called the adiabatic field and s is called the entropy field. The meaning of the names of the fields becomes clear when one considers their fluctuations.

The line-element for arbitrary scalar perturbations of the Robertson–Walker metric for a spatially flat universe reads (using the notation of [5])

$$ds^2 = -(1 + 2A)dt^2 + 2a^2 B_{,i} dx^i dt + a^2 [(1 - 2\psi)\delta_{ij} + 2E_{,ij}] dx^i dx^j. \quad (8)$$

The gauge-invariant curvature perturbation, defined as

$$\mathcal{R} = \psi + \frac{H\delta\rho}{\dot{\rho}}, \quad (9)$$

is, on very large scales, constant for purely adiabatic perturbations (see e.g. [14] and references therein). However, entropy perturbations are a source for the curvature perturbation (9). In addition, the entropy perturbation between two species A and B , defined as

$$\mathcal{S} = \frac{\delta n_A}{n_A} - \frac{\delta n_B}{n_B}, \quad (10)$$

where n_i are the number densities of the particle species i , can evolve in time, even on superhorizon scales. Therefore it was argued [13], that on very large scales in general we have the following equations describing the evolution of $\mathcal{R}$ and $\mathcal{S}$:

$$\dot{\mathcal{R}} = \alpha H \mathcal{S} \quad (11)$$

$$\dot{\mathcal{S}} = \beta H \mathcal{S} \quad (12)$$

For the case of the two slow-rolling scalar fields and in the spatial flat gauge ($\psi = 0$), $\mathcal{R}$ and $\mathcal{S}$ are given by

$$\mathcal{R} \approx \frac{H(\dot{\phi}\delta\phi + \dot{\chi}\delta\chi)}{\dot{\phi}^2 + \dot{\chi}^2} = \frac{H\delta\sigma}{\dot{\sigma}} \quad (13)$$

and

$$\mathcal{S} = \frac{H(\dot{\phi}\delta\chi - \dot{\chi}\delta\phi)}{\dot{\phi}^2 + \dot{\chi}^2} = \frac{H\delta s}{\dot{\sigma}} \quad (14)$$

Fluctuations in the field σ are adiabatic perturbations, whereas fluctuations in s are entropic perturbations. On very large scales ($k \ll aH$) and in the flat gauge the evolution of fluctuations are described as [5]:

$$(\delta\sigma)^{\cdot\cdot} + 3H(\delta\sigma)^{\cdot} + (V_{\sigma\sigma} - \dot{\theta}^2)\delta\sigma = -2V_{\sigma}A + \dot{\sigma}\dot{A} + 2(\dot{\theta}\delta s)^{\cdot} - 2\frac{V_{\sigma}}{\dot{\sigma}}\dot{\theta}\delta s \quad (15)$$

and

$$(\delta s)^{\cdot\cdot} + 3H(\delta s)^{\cdot} + (V_{ss} - \dot{\theta}^2)\delta s = -2\frac{\dot{\theta}}{\dot{\sigma}}[\dot{\sigma}((\delta\sigma)^{\cdot} - \dot{\sigma}A) - \ddot{\sigma}\delta\sigma]. \quad (16)$$

The metric perturbation A can be obtained from Einstein's equation and is given, in the flat gauge, by

$$HA = 4\pi G(\dot{\phi}\delta\phi + \dot{\chi}\delta\chi) \quad (17)$$

It was shown in [13] that for the case of two slow-rolling scalar fields, α and β in (11) and (12) are given in terms of the slow roll parameters

$$\alpha = -2\eta_{\sigma\sigma}, \quad (18)$$

$$\beta = -2\epsilon + \eta_{\sigma\sigma} - \eta_{ss}, \quad (19)$$

and are, therefore, specified by the potential $V(\phi, \chi)$. In the last two equations, the slow roll parameters are constructed from the usual slow-roll parameter for ϕ and χ (4) and are given by

$$\epsilon = \frac{1}{16\pi G} \left(\frac{V_{\sigma}}{V} \right)^2 \approx \epsilon_{\phi} + \epsilon_{\chi}. \quad (20)$$

and

$$\begin{aligned} \eta_{\sigma\sigma} &= \eta_{\phi\phi} \cos^2 \theta + 2\eta_{\phi\chi} \cos \theta \sin \theta + \eta_{\chi\chi} \sin^2 \theta, \\ \eta_{ss} &= \eta_{\phi\phi} \sin^2 \theta - 2\eta_{\phi\chi} \cos \theta \sin \theta + \eta_{\chi\chi} \cos^2 \theta, \\ \eta_{\sigma s} &= (\eta_{\chi\chi} - \eta_{\phi\phi}) \sin \theta \cos \theta + \eta_{\phi\chi} (\cos^2 \theta - \sin^2 \theta). \end{aligned} \quad (21)$$

The time evolution of $\mathcal{R}$ and $\mathcal{S}$ between horizon crossing and some later time are given by

$$\begin{pmatrix} \mathcal{R} \\ \mathcal{S} \end{pmatrix} = \begin{pmatrix} 1 & T_{\mathcal{R}\mathcal{S}} \\ 0 & T_{\mathcal{S}\mathcal{S}} \end{pmatrix} \begin{pmatrix} \mathcal{R} \\ \mathcal{S} \end{pmatrix}_*, \quad (22)$$

where the asterisk marks the time of horizon crossing. The transfer functions $T_{\mathcal{R}\mathcal{S}}$ and $T_{\mathcal{S}\mathcal{S}}$ are given by

$$\begin{aligned} T_{\mathcal{S}\mathcal{S}}(t, t_*) &= \exp \left(\int_{t_*}^t \beta(t') H(t') dt' \right) \\ T_{\mathcal{R}\mathcal{S}}(t, t_*) &= \int_{t_*}^t \alpha(t') T_{\mathcal{S}\mathcal{S}}(t_*, t') H(t') dt'. \end{aligned} \quad (23)$$

Thus, in the case of two slow rolling scalar fields, the transfer functions are completely specified by the potential through the slow-roll parameter. In the following we would like to find the changes of the evolution of adiabatic and entropy perturbations in brane world theories.

Brane World in AdS: In the case of a brane embedded in an Anti-de Sitter spacetime, Einstein's equation is not valid at high energies. Here, modifications become important in the form of a term which is quadratic in $T_{\mu\nu}$, the energy-momentum tensor of matter confined on the brane. In addition, the gravitational field in the bulk is encoded in the projection of the five-dimensional Weyl-tensor onto the brane. It was shown that the Einstein-equation has the form [15]

$$G_{\mu\nu} = \kappa_4^2 T_{\mu\nu} + \kappa_5^2 \Pi_{\mu\nu} - E_{\mu\nu}, \quad (24)$$

where $\Pi_{\mu\nu}$ is the term quadratic in $T_{\mu\nu}$ and $E_{\mu\nu}$ is the projected Weyl tensor. For the case of a homogeneous

and isotropic expanding brane embedded in an Anti-de Sitter space ($E_{\mu\nu} = 0$), the Friedmann equation is given by

$$H^2 = \frac{8\pi G}{3}\rho \left[1 + \frac{\rho}{2\lambda}\right]. \quad (25)$$

In this equation λ is the (intrinsic) brane tension and ρ is the total energy density of the scalar fields¹. The equations of motion for scalar fields confined on the brane are given by the usual Klein–Gordon equations (2) and (3).

It was shown in [9] that slow roll inflation driven by a single scalar field demands that the two slow-roll parameters

$$\epsilon \equiv \frac{1}{16\pi G} \left(\frac{V'}{V}\right)^2 \left[\frac{4\lambda(\lambda + V)}{(2\lambda + V)^2}\right] \quad (26)$$

$$\eta \equiv \frac{1}{8\pi G} \left(\frac{V''}{V}\right) \left[\frac{2\lambda}{2\lambda + V}\right] \quad (27)$$

are small. The number of e-folds is enhanced for a given potential compared to the usual four-dimensional case. It was also shown that the amplitude of scalar perturbations is enhanced relative to the standard result in four dimensions for a given potential.

In the case of the two scalar fields ϕ and χ , the set of slow-roll parameters is given by

$$\begin{aligned} \epsilon_\phi &\equiv \frac{1}{16\pi G} \left(\frac{V_\phi}{V}\right)^2 \left[\frac{4\lambda(\lambda + V)}{(2\lambda + V)^2}\right], \\ \epsilon_\chi &\equiv \frac{1}{16\pi G} \left(\frac{V_\chi}{V}\right)^2 \left[\frac{4\lambda(\lambda + V)}{(2\lambda + V)^2}\right], \\ \eta_{\phi\phi} &\equiv \frac{1}{8\pi G} \left(\frac{V_{\phi\phi}}{V}\right) \left[\frac{2\lambda}{2\lambda + V}\right], \\ \eta_{\phi\chi} &\equiv \frac{1}{8\pi G} \left(\frac{V_{\phi\chi}}{V}\right) \left[\frac{2\lambda}{2\lambda + V}\right], \\ \eta_{\chi\chi} &\equiv \frac{1}{8\pi G} \left(\frac{V_{\chi\chi}}{V}\right) \left[\frac{2\lambda}{2\lambda + V}\right], \end{aligned} \quad (28)$$

which implies the corresponding changes for the slow-roll parameters for the adiabatic field σ and s . The changes in perturbations on brane worlds are two-fold. First, we perturb the spacetime away from Anti-de Sitter, meaning that the projected Weyl-tensor no longer vanishes. i.e. $\delta E_{\mu\nu} \neq 0$. However, in the case of a de Sitter brane and on large scales, it was shown that the Weyl contribution is negligible (e.g. [16]). We therefore neglect that term. The second contribution is quadratic in $T_{\mu\nu}$, which we do not neglect. It implies that the metric perturbation A is now given by [17]

$$HA = 4\pi G \left(1 + \frac{\rho}{\lambda}\right) (\dot{\phi}\delta\phi + \dot{\chi}\delta\chi). \quad (29)$$

Using (15) and (16), it is not difficult to show that these changes imply that

$$H^{-1}(\delta\sigma)^\cdot \simeq (2\epsilon - \eta_{\sigma\sigma})\delta\sigma - 2\eta_{\sigma s}\delta s \quad (30)$$

and

$$H^{-1}(\delta s)^\cdot \simeq -\eta_{ss}\delta s, \quad (31)$$

on large scales in the slow roll regime. These are exactly the same expressions found in [13] for General Relativity. This implies that the evolution of $\mathcal{R}$ and $\mathcal{S}$ are given by the expressions

$$\dot{\mathcal{R}} \simeq -2H\eta_{\sigma s}\mathcal{S} \quad (32)$$

$$\dot{\mathcal{S}} \simeq [-2\epsilon - \eta_{ss} + \eta_{\sigma\sigma}]H\mathcal{S} \quad (33)$$

Thus, we have found that the equations have the same form as in usual cosmology based on General Relativity. Therefore, the parameters α and β are the same as in eq. (18) and (19), but for a given potential they are reduced by a factor $\approx \lambda/V$ at high energies on the brane. At very high energies, i.e. $V \gg \lambda$, we have that the ratios $\dot{\mathcal{R}}/\mathcal{S}$ and $\dot{\mathcal{S}}/\mathcal{S}$ are suppressed by a factor $\approx \sqrt{\lambda/V}$. That means that the transfer functions (23) are significantly modified in that case. Indeed, for very large energies we have $T_{SS} \rightarrow 1$ and $T_{RS} \rightarrow 0$ as $V/\lambda \rightarrow \infty$, meaning that entropy perturbations have less influence on adiabatic perturbations and even if entropy perturbations are created, the final correlation between the adiabatic and entropy modes would be suppressed.

Our findings imply that, even in models with step potentials (see [10]) and multiple scalar fields, the production of entropy perturbations is suppressed, as long as the energies are much higher than the brane tension.

It is not difficult to understand the physical reason for the results obtained so far. From equation (15) and (16) we see that the angle θ merely has to vary in order for the fields s and σ to interact². For the case of the slow-rolling scalar fields on the brane, the equation of motion for the angle θ in the slow roll regime is given by

$$\dot{\theta} \simeq -\eta_{\sigma s}H. \quad (34)$$

This equation holds also for General Relativity (see [13]). For a given potential, this equation implies that, for brane world inflation at high energies, θ is almost constant, and therefore that the background trajectory will quickly follow a straight line (see figure).

For the extreme case $V/\lambda \rightarrow \infty$, the equation above implies that $\dot{\theta} \rightarrow 0$. In that case there is no efficient transfer of energy between s and σ . Instead, s evolves

¹We have assumed that the four-dimensional cosmological constant is negligible.

²If θ is time varying, the background trajectory is curved.

like a background field which does not influence perturbations in σ and the gravitational potential. Note that this also means that any transfer of non-Gaussianity from the entropy field s to the gravitational potential and/or the adiabatic field σ is strongly suppressed (see [18] for a recent discussion).




FIG. 1. Schematic sketch of the evolution of ϕ and χ : the enhanced friction in brane worlds causes the fields quickly to move on a straight line after a short period of damping of the motion of the angle θ .

Note that our findings do not imply that there are no entropy perturbations generated in brane worlds. In this letter we have discussed only the evolution of perturbations *during* inflation. In principle the power spectra in the radiation dominated epoch depend on the details of the decay of the fields ϕ and χ and therefore a calculation has to be made using a detailed theory of preheating in two-field inflation.

After inflation, there is another source of entropy perturbation, namely the projected Weyl-tensor in equation (24), which acts as a source term for perturbations on the brane. It could, in principle, produce entropy perturbations even at late times [17]. However, the amplitude of the source term in (17) for linear perturbations around a homogeneous-isotropic brane universe has not been calculated so far.

To conclude, we have shown that the modifications to General Relativity implied by brane world scenarios can suppress the production of entropy perturbations during inflation driven by multiple scalar fields. The reason is that, for a given potential, the background trajectories are less curved in brane worlds due to the larger friction acting on the fields. In addition, if inflation is driven at energies much larger than the brane tension, the final power spectra do not depend on the initial conditions and/or trajectories of the fields. This also implies that any transfer of non-Gaussianity from the entropy field to the gravitational potential and/or adiabatic perturbations is suppressed. Thus, if any entropy perturbations are detected in the CMB and/or large scale structure, it would mean that either inflation on a brane happened at energies smaller than the brane tension or that steeper potentials have driven inflation. Although our calcula-

tions are based on the Randall-Sundrum model, we believe that our conclusions carry over to other brane world models as well, such as models with a bulk scalar field [19]. In these models, the brane tension is not necessarily constant but depends on the evolution of the bulk scalar field near the brane. As long as the time-evolution of the bulk scalar field is negligible compared to the evolution of the inflaton fields, our conclusions should hold. Otherwise, there will be a source for the curvature perturbation, coming from the apparent non-energy-conservation on the brane due to the bulk scalar field [20].

Acknowledgements: We are grateful for comments from C. Gordon and G. Rigopoulos. The authors are supported in part by PPARC.

-
- [1] A.R. Liddle, D. Lyth, *Cosmological Inflation and Large Scale Structure*, Cambridge University Press (2000)
 - [2] J. Garcia-Bellido, D. Wands, Phys.Rev. D **53**, 5437 (1996)
 - [3] R. Trotta, A. Riazuelo, R. Durrer, Phys. Rev. Lett. **87**, 231301 (2001)
 - [4] L. Amendola, C. Gordon, D. Wands, M. Sasaki, Phys. Rev. Lett. **88**, 211302 (2002)
 - [5] C. Gordon, D. Wands, B.A. Bassett, R. Maartens, Phys.Rev.D **63**, 023506 (2001)
 - [6] S. Groot Nibbelink, B.J.W. van Tent, Class.Quant.Grav. **19**, 613 (2002)
 - [7] N. Bartolo, S. Matarrese, A. Riotto, Phys.Rev. D **64**, 123504 (2001)
 - [8] L. Randall, R. Sundrum, Phys.Rev.Lett. **83**, 4690 (1999)
 - [9] R. Maartens, D. Wands, B.A. Bassett, I. Heard, Phys.Rev.D **62**, 041301 (2000)
 - [10] E.J. Copeland, A.R. Liddle, J.E. Lidsey, Phys.Rev. D **64**, 023509 (2001)
 - [11] R. Hawkins, J. Lidsey, Phys.Rev. D **63**, 041301 (2001)
 - [12] G. Huey, J. Lidsey, Phys.Lett.B **514**, 217 (2001); J. Lidsey, T. Matos, L. Urena-Lopez, Phys.Rev. D **66**, 023514 (2002)
 - [13] D. Wands, N. Bartolo, S. Matarrese, A. Riotto, astro-ph/0205253, to appear in Phys. Rev. D (2002)
 - [14] D. Wands, K. Malik, D. Lyth, A.R. Liddle, Phys.Rev. D **62**, 043527 (2000)
 - [15] T. Shiromizu, K. Maeda, M. Sasaki, Phys.Rev. D **62**, 024012 (2000)
 - [16] R. Maartens, Phys. Rev.D **62**, 084023 (2000); G. Giudice, E. Kolb, J. Lesgourgues, A. Riotto, hep-ph/0207145 (2002)
 - [17] D. Langlois, R. Maartens, M. Sasaki, D. Wands, Phys.Rev. D **63**, 084009 (2001)
 - [18] F. Bernardeau, J.-P. Uzan, hep-ph/0207295 (2002)
 - [19] A. Lukas, B.A. Ovrut, K.S. Stelle, D. Waldram, Phys.Rev. D **59**, 086001 (1999)
 - [20] C. van de Bruck, M. Dorca, C.J.A.P. Martins, M. Parry, Phys. Lett. B **495**, 183 (2000)